

Verlag Dr. Friedrich Pfeil
ISSN 0936-9902

Ichthyological Exploration of Freshwaters

An international journal for field-orientated ichthyology

**Volume 26
Number 3**



Ichthyological Exploration of Freshwaters

An international journal for field-orientated ichthyology

Volume 26 · Number 3 · November 2015
pages 193–288, 44 figs., 17 tabs.

Managing Editor

Maurice Kottelat, Rue des Rauragues 6
CH-2800 Delémont, Switzerland
Tel. +41 32 4623175 · Fax +41 32 4622259 · E-mail mkottelat@dplanet.ch

Associate Editors

Ralf Britz, Department of Zoology, The Natural History Museum, London, United Kingdom
Kevin W. Conway, Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, USA

Editorial board

Sven O. Kullander, Naturhistoriska Riksmuseet, Stockholm, Sweden
Helen K. Larson, Museum and Art Gallery of the Northern Territory, Darwin, Australia
Lukas Rüber, Naturhistorisches Museum, Bern, Switzerland
Ivan Sazima, Museu de Zoologia, Unicamp, Campinas, Brazil
Paul H. Skelton, South African Institute for Aquatic Biodiversity, Grahamstown, South Africa
Tan Heok Hui, Lee Kong Chian Natural History Museum, National University of Singapore, Singapore

Ichthyological Exploration of Freshwaters is published quarterly

Subscriptions should be addressed to the Publisher:

Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, 81379 München, Germany
PERSONAL SUBSCRIPTION : EURO 100 per Year/volume - 4 issues (includes surface mail shipping)
INSTITUTIONAL SUBSCRIPTION : EURO 180 per Year/volume - 4 issues (includes surface mail shipping)

Manuscripts should be addressed to the Managing Editor:
Maurice Kottelat, Rue des Rauragues 6, CH-2800 Delémont, Switzerland

CIP-Titelaufnahme der Deutschen Bibliothek

Ichthyological exploration of freshwaters : an international
journal for field-orientated ichthyology. – München : Pfeil.
Erscheint jährl. viermal. – Aufnahme nach Vol. 1, No. 1 (1990)
ISSN 0936-9902

Vol. 1, No. 1 (1990) –

Copyright © 2015 by Verlag Dr. Friedrich Pfeil, München, Germany

All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying or otherwise, without the prior permission of the copyright owner. Applications for such permission, with a statement of the purpose and extent of the reproduction, should be addressed to the Publisher, Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, 81379 München, Germany.

Printed by PBTisk a.s., Příbram I – Balonka

ISSN 0936-9902
Printed in the European Union

Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, 81379 München, Germany
Phone +49 89 742827-0 · Fax +49 89 7242772 · E-mail: info@pfeil-verlag.de · www.pfeil-verlag.de

The microhabitat, behavior and diet of *Centromochlus meridionalis*, a small catfish of Amazon streams (Teleostei: Auchenipteridae)

Fernando Gonçalves Cabeceira*, Denise Caragnato Parisotto**, Jansen Zuanon***,
 Alberto Akama**** and Lucélia Nobre Carvalho*,**

Centromochlus meridionalis was recently described from streams of Brazilian Amazon. The marbled dark brown and black color pattern together with the use of submersed leaf litter accumulations for shelter differentiates this species among its congeners. We present information about the biology of *C. meridionalis* based on data gathered in streams and under captivity, as well as on a dietary analysis. Behavioral observations were made in captivity (aquaria). Dietary analysis was performed based on stomach contents of 38 specimens. The species was found in 8 out of 12 small streams sampled, where specimens were captured predominantly amidst submerged leaf litter. Nine types of behavioral acts were identified, of which “swimming near the substratum” and “charging the substratum for food” were the most frequent. Thirty (~79 %) out of the 38 stomachs had food, and the diet was composed of 27 types of food items. *Centromochlus meridionalis* can be considered a generalist microcarnivore, consuming predominantly autochthonous and allochthonous insects, and other terrestrial invertebrates, crustaceans and fish as well. The use of different water column strata during foraging and the diversity of food items consumed indicate that this small catfish utilizes several feeding tactics.

Introduction

The catfish family Auchenipteridae is endemic of the Neotropics and broadly distributed in South America, with 20 genera and approximately 90

species (Ferraris, 2003, 2007). Most of those species are active at night and at dusk, seeking refuge during the day in deep water or in cavities in logs and rocks (Rodriguez et al., 1990). Auchenipteridae are the only siluriforms that possess a copulatory

* Programa de Pós-Graduação em Ecologia e Conservação da Biodiversidade, Universidade Federal de Mato Grosso – Câmpus Universitário de Cuiabá, Avenida Fernando Correa da Costa N° 2.367, Boa Esperança, 78060-900, Brazil. E-mail: fernando.cabeceira@gmail.com

** Universidade Federal de Mato Grosso, Instituto de Ciências Naturais, Humanas e Sociais, Câmpus Universitário de Sinop, Avenida Alexandre Ferronato N° 1.200, Setor Industrial, 78557-267, Brazil. E-mail: carvalholn@yahoo.com.br

*** Instituto Nacional de Pesquisas da Amazônia – INPA – Coordenação de Biodiversidade. Avenida André Araújo, n° 2936, Aleixo, 69060-000, Brazil. E-mail: jzuanon3@gmail.com

**** Museu Paraense Emílio Goeldi, Campus de Pesquisa, Coordenação de Zoologia, Avenida Magalhães Barata, 376, São Braz, 66040-170, Brazil. E-mail: aakama@gmail.com

structure and a reproductive pattern in which the male uses its modified anal fin to inseminate the female (Loir et al., 1989; Soares-Porto, 1998; Downing Meisner et al. 2000). Males of some species can also develop temporary structures, such as bony whiskers and hypertrophied fin spikes, during the reproductive period (Burgess, 1989; Akama & Ferraris, 2003). The Auchenipteridae is composed of two subfamilies: Auchenipterinae, which includes larger species that usually inhabit rivers, lakes and swamps, and Centromochlinae, including small species found in rivers as well as in small streams (Santos et al., 2004; Cabalzar et al., 2005).

Currently, the Centromochlinae includes 31 valid species, in the genera *Tatia*, *Glanidium*, *Gelanoglanis* and *Centromochlus* (Ferraris, 2007). Most of the literature on the subfamily involves the description of new species (Sarmiento-Soares & Buckup, 2005; Sarmiento-Soares & Martins-Pinheiro, 2008), a systematic review (Sarmiento-Soares & Martins-Pinheiro, 2008) and lists of species (Ferraris, 2003, 2007), with relatively little information on the biology of the species and related to feeding habits (Ortêncio Filho et al., 2001; Cassemiro et al., 2005). *Centromochlus* comprises 14 valid species (Brindelli et al., 2015). Most of the information about the biology of the genus is scattered in studies related to ichthyofaunal surveys (Mees, 1974; Santos et al., 2004; Souza et al., 2012), phylogenetic relationships (Birindelli, 2010), and a report of accidents caused by the spines (Sazima et al., 2005). All species of the genus (except *C. meridionalis*) have crepuscular habits, foraging near surface.

Centromochlus meridionalis is a small species inhabiting streams of the Teles Pires River basin, Tapajós basin (Sarmiento-Soares et al., 2013). Specimens were captured from underwater leaf litter accumulations in small 1st- and 2nd-order streams. In contrast, some congeners (e. g., *C. heckelii*) tend to occupy higher-order streams and large rivers (Sarmiento-Soares et al., 2013). *Centromochlus meridionalis* is similar to species of *Trachelyopterus* (Auchenipterinae) with respect to general morphology and especially color pattern (Sarmiento-Soares et al., 2013). The marbled dark brown and black body pattern may serve as cryptic coloration among the submersed leaves (FGC, pers. obs.). Therefore, the particular color pattern and the use of a microhabitat that is unusual for *Centromochlus* species (submerged leaf litter accumulations) indicate that this species may have

different behavioral characteristics from the other members of the genus. In this study, we present information about the behavior of *C. meridionalis* based on information about its natural habitat, its behavioral repertoire in captivity and the results of a dietary analysis of preserved specimens.

Materials and methods

Specimens of *Centromochlus meridionalis* were collected in 1st- and 2nd-order streams (cf. Strahler, 1957) of the Teles Pires River basin (11°35'59" S 55°15'21.0" W) in June and July 2011, during the dry season in the region. The streams are located in Amazon Forest remnants in the municipality of Cláudia, Mato Grosso, southern Brazilian Amazon. This region is covered by vegetation classified as Transitional Amazon Forest (Ferreira et al., 1999).

The fish were collected actively during the day by two collectors that explored all microhabitats in a stretch of 150 meters with sieves and hand nets. All the leaf litter in the stretch were searched for fish. Minnow traps were set in the same stretch in the afternoon of the day before and left overnight in the stream. The fish were kept alive in plastic boxes with aeration and transported to the laboratory of the Acervo Biológico da Amazônia Meridional (ABAM) at Universidade Federal de Mato Grosso (UFMT), Sinop campus.

Voucher specimens were deposited in: ABAM, fish collection of Acervo Biológico da Amazônia Meridional, Sinop; INPA, Instituto Nacional de Pesquisas da Amazônia, Manaus; MBML, Museu de Biologia Prof. Mello Leitão, Santa Teresa; and MNRJ, Museu Nacional do Rio de Janeiro, Rio de Janeiro. All from Brazil: Mato Grosso state: Loanda stream, a small tributary of Roquete river: ABAM I-00316, 12, 23.8–42.8 mm SL; 11°25'33.1" S 55°16'39.3" W; F. G. Cabeceira, W. S. de Moraes & J. Dambroz 24 July 2010. – ABAM I-00317, 11, 31.7–43.7 mm SL; 11°25'42.7" S 55°16'34.6" W; F. G. Cabeceira, W. S. de Moraes & Amanda F. Mortati, 25 July 2010. – ABAM I-00318, 9, 9.4–48.5 mm SL; 11°35'59.1" S 55°15'21.0" W; F. G. Cabeceira, W. S. de Moraes & Márcia C. V. dos Santos, 9 July 2010. – ABAM I-00398, 2, 53.2–60.2 mm SL; 11°25'48.7" S 55°20'16.30" W; F. G. Cabeceira & E. Barbosa, 6 May 2011. – INPA 37893, 10, 40.4–47.9 mm SL; 11°25'33.1" S 55°16'39.3" W; F. G. Cabeceira & E. Barbosa, 8 May 2011. – MBML 5617, 3, 32.2–46.2 mm SL; 11°25'33.1" S 55°16'39.3" W; F. G. Ca-

beceira & E. Barbosa, 8 May 2011. – MNRJ 40701, 2, 51.8–52.3 mm SL; 11°25'33.1" S 55°16'39.3" W; F. G. Cabeceira, W. S. de Moraes & J. Dambroz, 24 July 2010.

Habitat. The characterization of the habitat was based on descriptive parameters of the stream environment and on information about the structure of the submersed leaf accumulations. Environmental parameters were measured according to the protocol proposed by Mendonça et al. (2005). The stream's width, depth, current speed, vegetation (canopy) cover, and limnological data (pH, temperature) were measured. In meandering streams leaf litter accumulations occur in depositional areas, forming discrete (i. e. discontinuous) habitats to the aquatic fauna.

Behavioral observations. All the observations were performed on two groups of fish, each consisting of four specimens and housed in separate tanks (80 and 50 liters), both containing submerged branches and leaves on a sand substrate. The fish were fed with granulated commercial aquarium fish food and living coleopteran (Tenebrionidae) larvae. The behavioral observations were performed only in the larger tank. During the observation sessions, a blind (a black cloth) separated the tank from the observer to avoid disturbing the fish. After 10 observation sessions, the fish group in the larger tank was changed and a similar set of observations was made with the second group.

Before conducting the quantitative behavioral measurements, 16 hours of preliminary observations were made, aiming to identify the main activity period and to record the behavioral repertoire. To quantify the observed behavioral acts, standardized observation sessions were conducted at night from 6:00 pm to 6:00 am, divided into 13 sessions of 20 minutes each and totaling 4.3 hours. Each group of *C. meridionalis* was observed for 10 non-consecutive nights, resulting in a total of 86 hours of observations for both groups. The observations were performed using all-occurrences sampling for a general characterization of the behavior and focal-animal sampling to count the number of times a given fish performed each behavior (Altman, 1974). The results are presented as frequencies of occurrence (%) in relation to the total behavioral acts recorded.

Diet analysis. The diet was analyzed based on the stomach contents of 38 specimens (ABAM I-00316, ABAM I-00317, ABAM I-00318) preserved immediately after collecting in the field. The fish were measured (standard length, SL, in mm) and weighed (g), and the stomach contents were analyzed under a stereoscopic magnifier. The food items were identified to the lowest taxonomic level possible and later grouped into seven categories: fish, crustaceans, aquatic insects, terrestrial insects, other terrestrial arthropods, insect fragments and organic debris.

The fullness level of each stomach was estimated visually and categorized as follows: 0 % = empty stomach; 10 % = 0–10 % of the volume of the stomach filled with food; 25 % = 10–25 %; 50 % = 25–50 %; 75 % = 50–75 %; and 100 % = stomach totally full (adapted from Goulding et al., 1988). The relative volume (R.V.%) occupied by each item in the stomach was assessed considering the whole amount of food in the stomach as 100 %, according to Soares (1979). To correct the R.V.% values for the differences in the amount of food in each stomach, the values of relative abundance were then multiplied by the degree of stomach fullness. Finally, the R.V.% values were calculated by dividing the sum of values for each food category by the total volume of food consumed by the specimens analyzed. The frequency of occurrence (F.O.%) of the food items was also assessed as proposed by Hyslop (1980), according to the formula $F.O.\% = (n^\circ \text{ of stomachs with item } i / n^\circ \text{ of stomachs with food}) \times 100$. These two parameters (R.V.% and F.O.%) were used to calculate the Alimentary Index (A.I.) (Kawakami & Vazzoler, 1980), in which $A.I. = (R.V.\% \times F.O.\% / \sum R.V.\% \times F.O.\%) \times 100$.

Results

Microhabitat. Specimens of *C. meridionalis* (Fig. 1) were collected in 1st- and 2nd-order streams 1.8–3.2 m wide and 0.2–0.5 m deep, current speed 0.2–0.4 m · s⁻¹, and discharge 0.1–0.3 m³ · s⁻¹. Vegetation cover by riparian forest canopy averaged 88.6 %. The water was acidic (mean pH 4.9; min = 4.2, max = 5.4), and the mean temperature was 23.7 °C (min = 22.8, max = 24.5). The stream bed was composed primarily of sand (45 %; grains approximately 0.6–2.0 mm in diameter), fine sediment (20.4 %), coarse litter (17.6 %; leaves and small branches), fine litter (8 %; finely particulate




Fig. 1. *Centromochlus meridionalis* photographed alive in aquarium. **a**, female, 60.2 mm SL; and **b**, male, 53.2 mm SL (ABAM I-00398). Arrow indicates modified anal fin forming the intromittent organ. (Photographs L. N. Carvalho and F. G. Cabeceira).

plant matter) and logs (4.4 %; wood with diameter greater than 10 cm).

Centromochlus meridionalis was found in 8 of 12 streams sampled in the Teles Pires River basin, and co-occurred with 30 other fish species (FGC, pers. obs.). In a survey performed with standardized sampling effort during the dry season of 2010–2011, *C. meridionalis* was the second most abundant fish species in the studied streams, outnumbered only by *Moenkhausia phaeonota* (Characiformes: Characidae). An average of 9 individuals (range: 1–32) of *C. meridionalis* were collected per 150-m long stream stretch. The cat-

fish was primarily captured in loosely submersed leaf litter accumulations. On most of those occasions, two or three specimens were captured in a single leaf litter pack, with a maximum of five individuals (in that case, three adults and two juveniles). Individuals were also captured under submerged logs greater than 10 cm in diameter.

Behavior in aquarium. Behavioral observations revealed peak activity between 8:00 pm and 3:00 am, with individuals resting after dawn. Nine distinct behavioral acts were identified and divided into four categories: resting, moving,

Table 1. Characterization and frequency of behavioral acts performed by individuals of *Centromochlus meridionalis* in captivity. Total number of behavioral acts = 4190.

Behavioral category Behavioral act	Description	Frequency
Resting		
Resting over sand	Remains exposed over sand substrate	5.9 %
Resting among leaves and logs	Rests between submerged leaves and logs	13.3 %
Settled on the substrate	Undulating body movements while stationary, sometimes disrupting sand particles and dead leaves	1.6 %
Movement		
Swimming near the substrate	Swimming close to the substrate (maximum 5 cm above it)	45.6 %
Swimming in midwater	Swimming in middle layer of water column (5 cm above substrate and 5 cm below surface)	7.7 %
Swimming close to the surface	Swimming in top 5 cm of water column	6.9 %
Feeding		
Swimming rapidly toward food on the substrate	During swimming close to the substrate and after detecting food, the individual assumes a head-down position and swims forward rapidly, biting the food item	17.5 %
Picking food at the surface	The individual swims rapidly towards food floating at the surface, engulf it and returns rapidly to midwater	1.4 %
Interaction		
Agonistic interactions	When two individuals moving around the tank approach each other closely (less than 2 cm), both suddenly move away to avoid physical contact	0.1 %




Fig. 2. Illustration of behavioral acts performed by *Centromochlus meridionalis*: **a**, resting between leaves; and **b**, feeding at the bottom in head-down posture. For a description of the behaviors see Table 1. (Illustrations by Cristiane C. de Freitas).

feeding and agonistic interactions (Table 1; Fig. 2). The most frequent behavior was swimming near the substrate (Frequency of Occurrence = 34 %), food searching in the substrate (17.5 %), resting among leaves and logs (13.3 %) and swimming among submerged leaves and logs (11.6 %).

Movements across the aquarium were performed by rapid swimming and sudden changes in direction. During swimming, the fish often remained close to the substrate but also explored the middle layer of the water column at times. Swimming near the surface was less frequent and always occurred for very short periods (a few seconds) before returning to lower layers. At times, the fish quickly swam to the surface and returned to the bottom. The greatest activity was observed during feeding. When food was offered, the fish swam frenetically close to the substrate throughout the tank, over open sand and between leaves and logs. After locating a food particle, the catfish quickly positioned head-down and bit the food, using rapid mouth movements (Fig. 2b). Only during the last observation sessions the fish began to search for food on the water surface, which was interpreted as conditioned feeding behavior.

Diet. The specimens analyzed for stomach content analysis measured 9.3–52 mm SL and weighed 0.1–4.2 g. Of the 38 stomachs analyzed, 8 (21.1 %) were empty, 8 (21.1 %) showed up to 10 % of fullness, 3 (7.9 %) were 25 % full, 5 (13.2 %) were

50 % full, 9 (23.7 %) were 75 % full, and 5 (13.2 %) were 100 % full. The food items were grouped in 13 categories (Table 2). In terms of volume, debris appear with R.V. = 25 %, Ephemeroptera with R.V. 16 %, Hymenoptera (ants) with R.V. 15 % and Trichoptera with R.V. = 15 %. The most frequent

Table 2. Values of Alimentary Index (A.I.%) for the items found in stomach contents of 30 specimens of *Centromochlus meridionalis*, with their respective Frequency of Occurrence (F.O.%) and Relative Volume (R.V.%).

Food item	F.O.%	R.V.%	A.I.%
Insect fragments	13	3	1.4
Terrestrial insects			
Hymenoptera (ants)	33	15	18.7
Terrestrial arthropods			
Spirostreptida (Diplopoda)	3	2	0.3
Aquatic insects (mostly larvae)			
Trichoptera	37	15	21.1
Ephemeroptera	10	16	6
Hemiptera	10	2	0.6
Plecoptera	3	2	0.2
Megaloptera	3	1	0.1
Diptera	3	0.2	0.02
Coleoptera	17	9	5.4
Fish			
<i>Moenkhausia phaeonota</i>	7	2	0.3
Crustaceans			
Decapoda (shrimps)	3	2	0.3
Debris	47	25	44

items were debris with F.O. = 47 %, Trichoptera with 37 %, and Hymenoptera (ants) with 33 %. In terms of Alimentary Index, the most important items were organic debris (A.I. = 40 %), Trichoptera larvae (A.I. = 21.1 %) and adult Hymenoptera (ants) (A.I. = 18.7 %).

Discussion

Centromochlus meridionalis was found in streams with structural and limnological features similar to those of the small Central Amazon streams (Espírito-Santo et al., 2009). However, the substrate of the studied streams was covered by a smaller amount of coarse litter than that present in Central Amazon streams (Mendonça et al., 2005; Carvalho, 2008). This apparent scarcity of coarse litter suggests that *C. meridionalis* present a strong selection for this type of substrate, which is used as shelter during the day (FGC, pers. obs.). Considering the distinctive mottled color pattern of *C. meridionalis*, we hypothesize that use of coarse litter packs as resting sites is part of its primary defense system, by providing cover and camouflage (Keenleyside, 1979), which could be an effective strategy against visually-oriented predators. Available data for other species of Centromochlinae indicate that most of them generally inhabit larger streams and rivers (e.g., Py-Daniel et al., 2007; our personal observations), where they use various types of substratum for resting, such as rubble, cracks and small cavities in boulders and logs (e.g., Barreto & Rocha, 2005; Sarmiento-Soares & Buckup, 2005; Py-Daniel et al., 2007). Therefore, the use of litter packs for shelter and foraging by *C. meridionalis* appears to represent unusual features for a member of this subfamily, more common in fishes of the subfamily Auchenipterinae.

The behavioral observations recorded in aquaria showed that *C. meridionalis* has nocturnal habits similar to the majority of species of Auchenipteridae (Burgess, 1989; Akama & Ferraris, 2003). In captivity, however, the individuals of this species did not show the dusk activity peak known for other Centromochlinae (e.g., Ferraris, 2003; Sazima et al., 2005). During the day or under artificial light, the fish hid under leaves and logs at the bottom of the tank. Thus, the observed behaviors show that the species uses predominantly the lower layer of the water column both during resting and activity periods. This behavioral pat-

tern contrasts to the typical behavior known for most species of Centromochlinae, which usually swim close to the surface during foraging, searching for insects that fall into the water (Ferraris, 2003). However, such behavior was not commonly observed for *C. meridionalis* in the aquarium. The relatively short and stout body of *C. meridionalis* is also indicative of its nektobenthic habits, a trait more commonly observed in species of Auchenipterinae.

Centromochlus meridionalis showed a distinctive feeding behavior, foraging on the bottom by biting at food items from a head-down position. This feeding tactic may be associated with the main types of food items ingested, which included both terrestrial and aquatic insects possibly dragged by the current close to the bottom. Siluriforms show a variety of predatory tactics depending on the environmental conditions and prey types (Pohlmann et al., 2004). *Centromochlus meridionalis* may be considered a generalized microcarnivore because its diet primarily includes items of animal origin, such as autochthonous and allochthonous insects, other terrestrial invertebrates, crustaceans and (occasionally) small fish. Other species of Centromochlinae, e.g., *Glanidium ribeiroi* (Ortêncio Filho et al., 2001) and *G. bockmanni* (Sarmiento-Soares & Buckup, 2005) feed primarily on autochthonous and allochthonous insects (Ferraris, 2003). Adult *G. ribeiroi* may also feed on fish and plant matter (Ortêncio Filho, 2001), whereas *G. albescens* is considered an omnivorous species (Araújo et al., 2009). Nevertheless, there is no detailed information about the feeding behavior of those catfishes, which denotes the need for more behavioral and naturalistic studies focusing on species of Centromochlinae.

Although a considerable amount of organic debris was found in the stomach contents of *C. meridionalis*, this finding does not necessarily indicate that the species is a detritivore. Debris may have been accidentally ingested along with other food items collected at the stream bottom. Approximately 20 % of the substrate in the streams where the species was collected consisted of fine sediments similar to the debris found in the stomach contents. Fine sediment accumulates in slow-flowing stretches and near the margins of streams, where most of the leaf litter accumulations inhabited by *C. meridionalis* is found. Moreover, they ingested a considerable amount of trichopteran larvae of the family Leptoceridae,

which live buried in the mud (Mugnai et al., 2010), which may help explain the debris found in the stomach and also the unusual head-down posture employed during foraging.

The use of different layers of the water column by *C. meridionalis* during foraging, as well as the diversity of food items ingested, indicate a generalized microcarnivore diet. The food items ingested by *C. meridionalis* included the characid *Moenkhausia phaeonota* and small shrimps. Although those items are not particularly representative of the diet of *C. meridionalis*, they suggest the use of varied foraging tactics. *Moenkhausia phaeonota* is a diurnal fish that is abundant in the studied streams. It swims rapidly in small mid-water schools (sensu Casatti et al., 2001; Padial et al., 2009; FGC, pers. obs.). This characid species rests close to the bottom and along the stream margins at night, where it remains stationary or moves very slowly (FGC, pers. obs.). Due to those differences in activity time and microhabitat use, *C. meridionalis* most likely captures small fish while the prey rests at night.

Finally, it is necessary to remark that the streams inhabited by *C. meridionalis* are located in an area that is strongly threatened by human-induced disturbances, mainly resulting from the deforestation for agricultural use. In this sense, the restricted geographic distribution of *C. meridionalis* emphasizes the extinction risk to the species and reinforces the need to establish conservation measures to protect the headwaters of the Teles Pires River in the upper Tapajós River Basin, a region known for its high number of endemic fish species.

Acknowledgments

Fieldwork was funded by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (processes 569382/2008-4 and 556858/2009-3). We thank Edivaldo Barbosa for help during the field work. We are also grateful to Continental Farm for allowing access to the streams. Permission for collecting specimens was granted by ICMBio to FGC. JZ receives a productivity grant from CNPq (307464/2014-1). FGC thanks Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for a scholarship grant. This is publication number 47 of the Technical Series of the Núcleo de Estudos da Amazônia Meridional (NEBAM), and number 38 of Projeto Igarapés.

Literature cited

- Altman, J. 1974. Observational study of behavior: sampling methods. *Behaviour*, 49: 227–267.
- Araújo, F. G., B. C. T. Pinto & T. P. Teixeira. 2009. Longitudinal patterns of fish assemblages in a large tropical river in southeastern Brazil: evaluating environmental influences and some concepts in river ecology. *Hydrobiologia*, 618: 89–107.
- Akama, A. & C. J. Ferraris. 2003. *Entomocorus melaphareus*, a new species of auchenipterid catfish (Osteichthyes: Siluriformes) from the lower and middle reaches of the rio Amazonas. *Neotropical Ichthyology*, 1: 77–82.
- Barreto A. P. & J. M. Rocha. 2005. Assembleia de peixes de um riacho da Floresta Atlântica: composição e distribuição espacial (Guaraqueçaba, Paraná, Brasil). *Acta Scientiarum Biological Science*, 27: 153–160.
- Birindelli, J. L. O. 2010. Relações filogenéticas da superfamília Doradoidea (Ostariophysi, Siluriformes). Unpublished Doctoral Tesis, Universidade de São Paulo, São Paulo, 376 pp.
- Birindelli, J. L. O., L. M. Sarmento-Soares & F. C. T. Lima. 2015. A new species of *Centromochlus* (Siluriformes, Auchenipteridae, Centromochlinae) from the middle Rio Tocantins basin, Brazil. *Journal of Fish Biology*, 87: 860–875.
- Burgess, W. E. 1989. An atlas of freshwater and marine catfishes: a preliminary survey of the Siluriformes. TFH Publications, Neptune, 784 pp.
- Cabalzar, A., F. C. T. Lima & M. C. Lopes. 2005. Peixe e Gente no Alto Rio Tiquié: conhecimento tukano e tuyuka, ictiologia, etnologia. São Paulo, Instituto Socioambiental, 339 pp.
- Casatti, L., F. Langeani & R. M. C. Castro. 2001. Peixes de Riacho do Parque Estadual Morro do Diabo, Bacia do Alto Rio Paraná, SP. *Biota Neotropica*, 1: 1–15.
- Casemiro, F. A. S., N. S. Hahn & R. L. Delariva. 2005. Estrutura trófica da ictiofauna, ao longo do gradiente longitudinal do reservatório de Salto Caxias (rio Iguaçu, Paraná, Brasil), no terceiro ano após o represamento. *Maringá*, 27: 63–71.
- Carvalho, L. N. 2008. História natural de peixes de igarapés amazônicos: utilizando a abordagem do Conceito do Rio Contínuo. Unpublished Doctoral Tesis, Instituto Nacional de Pesquisas da Amazônia, Manaus, 142 pp.
- Downing Meisner, A. D., J. R. Burns, S. H. Weitzman & L. R. Malabarba. 2000. Morphology and histology of the male reproductive system in two species of internally inseminating South American catfishes, *Trachelyopterus lucenai* and *T. galeatus* (Teleostei: Auchenipteridae). *Journal of Morphology*, 246: 131–141.
- Espírito-Santo, H. M. V., W. E. Magnusson, J. Zuanon, F. P. Mendonça & V. L. Landeiro. 2009. Reproductive strategies of Amazonian stream fishes and their

- fine-scale use of habitat are ordered along a hydrological gradient. *Freshwater Biology*, 54: 536–548.
- Ferraris, C. J. 2003. Auchenipteridae (driftwood catfishes). Pp. 470–482 in: R. E. Reis, S. O. Kullander & C. J. Ferraris (eds), Checklist of the freshwater fishes of South and Central America. Editora Universitária da Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre.
- 2007. Checklist of catfishes, recent and fossil (Osteichthyes: Siluriformes), and catalogue of siluriform primary types. *Zootaxa*, 1418: 1–628.
- Ferreira, L. V., R. L. Sá, R. Buschbacher, G. Batmanian, J. M. C. Silva & E. Moretti. 1999. Identificação de áreas prioritárias para a Conservação da Biodiversidade através da Representatividade das Unidades de Conservação e Tipos de Vegetação nas Ecorregiões da Amazônia Brasileira. Ações Prioritárias para a Conservação da Biodiversidade da Amazônia. Programa Nacional da Diversidade Biológica, PROBIO, Ministério do Meio Ambiente.
- Goulding, M., M. L. Carvalho & E. J. G. Ferreira. 1988. Rio Negro: rich life in poor water. SPB Academic Publishing, The Hague, 200 pp.
- Hyslop, E. J. 1980. Stomach contents analysis: a review of methods and their application. *Journal of Fish Biology*, 17: 411–429.
- Kawakami, E. & G. Vazzoler. 1980. Método gráfico e estimativa de índice alimentar aplicado no estudo de alimentação de peixes. *Boletim do Instituto Oceanográfico*, 29: 205–207.
- Keenleyside, M. H. 1979. Diversity and adaptation in fish behaviour. Springer, Berlin, 208 pp.
- Loir, M., C. Cauty, P. Planquette & P. Y. Bail. 1989. Comparative study of the male reproductive tract in seven families of South-American catfishes. *Aquatic Living Resource*, 2: 45–56.
- Mees, G. F. 1974. The Auchenipteridae and Pimelodidae of Suriname (Pisces, Nematognathi). *Zoologische Verhandelingen*, 132: 1–256, 15 pls.
- Mendonça, F. P., W. E. Magnusson & J. Zuanon. 2005. Relationships between habitat characteristics and fish assemblages in small streams of Central Amazonia. *Copeia*, 2005: 750–763.
- Mugnai, R., J. L. Nessimian & D. F. Baptista. 2010. Manual de identificação de macro invertebrados aquáticos do Estado do Rio de Janeiro. Technical Books, Rio de Janeiro, 176 pp.
- Ortêncio Filho, H., N. S. Hahn, R. Fugui, & M. R. Russo. 2001. Aspectos da alimentação de *Glanidium ribeiroi* (Haseman, 1911) (Teleostei, Auchenipteridae), espécie endêmica do rio Iguaçú, PR. *Acta Limnológica*, 13: 85–92.
- Padial, A. A., S. M. Thomaz & A. A. Agostinho. 2009. Effects of structural heterogeneity provided by the floating macrophyte *Eichhornia azurea* on the predation efficiency and habitat use of the small Neotropical fish *Moenkhausia sanctaefilomenae*. *Hydrobiologia*, 624: 161–170.
- Py-Daniel, L. H. R., C. P. Deus, O. M. Ribeiro & L. M. Sousa. 2007. Peixes. Pp. 89–125. In: L. H. Rapp Py-Daniel et al. (eds.), Biodiversidade do Médio Madeira: bases científicas para proposta de conservação. Ed 1. Ministério do Meio Ambiente: Ministério da Ciência e Tecnologia, Manaus.
- Pohlmann, K. J., J. Atema & T. Breithaupt. 2004. The importance of the lateral line in nocturnal predation of piscivorous catfish. *Journal of Experimental Biology*, 207: 2971–2978.
- Rodriguez, M. A., S. E. Richardson & W. M. Lewis. 1990. Nocturnal behavior and aspects of the ecology of a driftwood catfish, *Entomocorus gameroi* (Auchenipteridae). *Biotropica*, 22: 435–438.
- Santos, G. M., B. Mérona, A. Afonso & M. Jégu. 2004. Peixes do Baixo Rio Tocantins: 20 anos depois da Usina Hidrelétrica Tucuruí. Eletronorte, Brasília, 216 pp.
- Sarmento-Soares, L. M. & R. F. Martins-Pinheiro. 2008. A systematic revision of *Tatia* (Siluriformes: Auchenipteridae: Centromochlinae). *Neotropical Ichthyology*, 6: 495–542.
- Sarmento-Soares, L. M. & P. Buckup. 2005. A New *Glanidium* from the Rio São Francisco Basin, Brazil (Siluriformes: Auchenipteridae: Centromochlinae). *Copeia*, 2005: 846–853.
- Sarmento-Soares, L. M., F. Cabeceira, L. N. Carvalho, A. Akama & J. Zuanon. 2013. *Centromochlus meridionalis*, a new catfish species from the southern Amazonian limits, Mato Grosso, Brazil (Siluriformes, Auchenipteridae). *Neotropical Ichthyology*, 11: 797–808.
- Sazima, I., J. Zuanon & V. Haddad. 2005. Puncture wounds by driftwood catfish during bucket baths: local habits of riverside people and fish natural history in the Amazon. *Wilderness and Environmental Medicine*, 16: 204–208.
- Soares, M. G. M. 1979. Aspectos ecológicos (alimentação e reprodução) dos peixes do igarapé do Porto, Aripuanã, MT. *Acta Amazonica*, 9: 325–352.
- Soares-Porto, L. M. 1998. Monophyly and interrelationships of the Centromochlinae (Siluriformes, Auchenipteridae). Pp. 331–350 in: L. R. Malabarba et al. (eds.), Phylogeny and classification of neotropical fishes. Editora Universitária da Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, 603 pp.
- Souza, L. S., J. W. Armbruster & D. C. Wemeke. 2012. The influence of the Rupununi portal on distribution of freshwater fish in the Rupununi district, Guyana. *Cybio*, 36: 31–43.
- Strahler, A. N. 1957. Quantitative analysis of watershed geomorphology. *American Geophysical Union Transactions*, 38: 913–920.

Received 21 October 2014

Revised 29 May 2015

Accepted 12 August 2015

Ichthyological Exploration of Freshwaters

An international journal for field-orientated ichthyology

INSTRUCTIONS TO CONTRIBUTORS

Warning

Prospective authors should read carefully the following instructions and follow them when submitting a manuscript. Doing so significantly hastens publication and saves money and efforts. Manuscripts which do not satisfy the instructions below may be rejected at the Editor's discretion and will not be returned.

Submission of manuscripts

The original manuscript should be sent to the Editor, Maurice Kottelat, by e-mail (mkottelat@planet.ch). Additional information is requested:

- 1) the name, postal and e-mail addresses, telephone and fax numbers of the corresponding author;
- 2) the names, postal and e-mail addresses of up to four persons outside the authors' institutions who are qualified to review the paper; and
- 3) a statement that the material has not been published and is not considered for publication elsewhere and that it will not be submitted elsewhere unless it is rejected or withdrawn. In submitting a manuscript, the author(s) accept(s) transfer of the copyright to the Publisher.

Co-authors, corresponding author

Authors are those who have played a significant role in designing and conducting the research and in writing the manuscript. Individuals who have only collected data, provided material or financial support, or reviewed the manuscript should be listed in acknowledgments. Honorary authorship is not accepted.

Co-authors should designate a single corresponding author to whom correspondence and proofs will be sent. All correspondence regarding the paper should go through the corresponding author. Correspondence will not be sent to other co-authors and correspondence from other co-authors regarding the manuscript will neither be answered nor taken into consideration.

Format

Files. The manuscript should be submitted in DOC or RTF format only. The text, captions, tables etc. must all be included in the same file. If the manuscript includes only a few illustrations, include them in low resolution in the word file. If the manuscript includes numerous illustrations they must be submitted in a separate PDF file; send all figures in low resolution and with caption in a single file. The files should be less than 8 MB.

Text. All manuscripts are subject to editorial revision before final acceptance for publication. Nothing in the manuscript should be underlined. Titles with numerical series designations are not permitted. Titles should be brief, fewer than 20 words and should indicate clearly the field of study and the group of fishes investigated. All abbreviations should be explained in the Method section (or figure caption when appropriate) or a reference to published explanations should be provided; exceptions are very common abbreviations, such as mm, km, kg, sec, min, yr, vs., SL. Footnotes are not permitted. All measurements must be in metric units. The first page should include: title of the paper, author(s), addresses and abstract, all left justified. The text should be followed by Material Examined (if appropriate), Acknowledgments (if any), Appendix (if any) and Literature Cited, in that order. Keys are desirable in taxonomic papers. They should be dichotomous and not serially indented.

Nomenclature. Names of living organisms should follow the appropriate and current International Codes of Nomenclature. Only formal names of genera and species should be written in italics. Names of authors and publication dates of scientific names should be mentioned once, in introduction or discussion, depending where most convenient, exceptionally as a table; bibliographical references must be included in the Literature cited section. Very old and classical works can be omitted if not absolutely justified.

Language. Manuscripts should be written in English. All papers must have a concise but informative abstract in English. In taxonomic papers, the abstract must include at least clear diagnosis of the new taxa. This maybe omitted for papers including the descriptions of many new taxa; consult the editor first. A second abstract, provided by the author(s), in the language of the country or area concerned by the text is acceptable. A maximum of two abstracts is permitted.

Acknowledgments. Identify individuals by first name(s) and surname. Do not list titles, position or institution. Acknowledge individuals, not positions. Idiosyncrasy and private jokes are not permitted.

Literature cited. Format for Literature Cited is that of the most recent issue. Do not abbreviate the names of journals. For books, give full name of publishing company or institution, and city. Manuscripts in preparation, abstracts, in-house reports and other literature not obtainable through

normal library channels cannot be cited. In-press manuscripts can be cited only if they have been formally accepted.

Tables. Tables should be included in the text file, at the end. Use Word format and do not anchor them. Tables must be numbered sequentially with Arabic numerals; they should have concise but self-explanatory headings. Do not insert frames, vertical rules, dotted lines or footnotes. The location of first citation of each table should be clearly indicated in the text.

Figures. Detailed instructions for the preparation of digital images are here: <http://pfeil-verlag.de/div/eimtag.php>

For the submission of new manuscript only low resolution copies are needed. Do not send large files at this stage. Case by case, if needed, we may ask you to send the original files at the time of submission.

All maps, graphs, charts, drawings and photographs are regarded as figures and are to be numbered consecutively and in the sequence of their first citation in the text. When several charts or photographs are grouped as one figure, they must be trimmed and spaced as intended for final reproduction. Each part of such a group figure should be lettered with a lower case block letter in the lower left corner. Where needed, scale should be indicated on the figure by a scale bar.

All illustrations should be designed to fit a width of 68 or 140 mm and a depth no greater than 200 mm. Lettering should be large enough to be easily seen when reduced onto a journal column (68 mm).

If a vector-graphics program is used, the original files saved by this program and all linked files must be submitted. Do not export or save the figure in a different format (for more details see the informations on <http://pfeil-verlag.de/div/eimtag.php>

If line drawings are scanned, the resolution must be 1200 dpi or more and the format must be bitmap (1 pixel = 1 bit).

If halftones are scanned, the resolution should never be lower than 400 dpi, applied to a width of 14 cm, even for photographs designed for column width.

Photographic prints and slides and original drawings must be scanned for submission. We will ask to send the original after acceptance of the manuscript.

Colour illustrations should preferably be submitted as slides (photographic slides, not slides prepared by a printer). Digital images should be only unmodified (raw) data files as originally saved by the camera or the scanner. If the data files are modified, a copy of the original, unmodified file should be submitted too.

The decision to print in colour or in black and white any figure originally submitted in colour remains with the editor and publisher. This decision will be based on scientific justification, quality of the original, layout and other editorial, financial and production constraints. By submitting colour originals, the authors know and accept that they may be published in black and white.

Review

Each manuscript will be sent to two reviewers for confidential evaluation. When justified, the reviewer's comments will be forwarded to the corresponding author. When submitting a revised manuscript, authors should briefly indicate the reasons for disregarding any suggestion they consider unacceptable. Remember that if a reviewer had questions or did not understand you, other readers may make the same experience and the answers should be in the manuscript and not in a letter to the editor. Changes in style, format and layout requested by the Editor are non-negotiable and non-observance will result in rejection of the manuscript.

Revised manuscripts received more than 6 months after the reviewers' comments had been sent will not be considered or will be treated as new submissions.

Proofs, Reprints and Page Charges

A PDF proof file will be sent to the corresponding author; it should be checked and returned to the Editor within one week. If corrections are not received within this delay, they may be done by the Editor, at the author's risks. Authors may be charged for any changes other than printer's error. Reprint orders must be forwarded with the corrections. The corresponding author is responsible for contacting the co-authors and forwarding their reprint orders.

The authors will receive a PDF file for personal use free of charge; high-resolution PDF files for unlimited use may be ordered. There will be no page charges and no charges for justified colour illustrations.

Ichthyological Exploration of Freshwaters

An international journal for field-orientated ichthyology

Volume 26 • Number 3 • November 2015

CONTENTS

Pavanelli, Carla Simone and Wayne C. Starnes: Revision of the trans-Andean scrapetooths genus <i>Saccodon</i> (Ostariophysi: Characiformes: Parodontidae).....	193
Mattos, José L. O., Wilson J. E. M. Costa and Alexandre C. A. Santos: <i>Geophagus diamantinensis</i> , a new species of the <i>G. brasiliensis</i> species group from Chapada Diamantina, north-eastern Brazil (Cichlidae: Geophagini).....	209
Cabeceira, Fernando Gonçalves, Denise Caragnato Parisotto, Jansen Zuanon, Alberto Akama and Lucélia Nobre Carvalho: The microhabitat, behavior and diet of <i>Centromochlus meridionalis</i> , a small catfish of Amazon streams (Teleostei: Auchenipteridae).....	221
Ng, Heok Hee and Wansheng Jiang: Intrafamilial relationships of the Asian hillstream catfish family Sisoridae (Teleostei: Siluriformes) inferred from nuclear and mitochondrial DNA sequences	229
Valdesalici, Stefano: <i>Anablepsoides fransvermeuleni</i> , a new killifish from Suriname (Teleostei: Rivulidae).....	241
Du Li-Na, Lu Yu-Fa and Chen Xiao-Yong: <i>Yunnanilus qujinensis</i> , a new species of loach from Yunnan, southwest China (Teleostei: Nemacheilidae)	249
Bragança, Pedro H. N. de , Felipe P. Ottoni and Filipe S. Rangel-Pereira: <i>Hyphessobrycon ellisae</i> , a new species from northeastern Brazil (Teleostei: Characidae).....	255
Shangningam, Bungdon and Waikhom Vishwanath: Two new species of <i>Garra</i> from the Chindwin basin, India (Teleostei: Cyprinidae).....	263
Decru, Eva, Jos Snoeks and Emmanuel Vreven: Taxonomic evaluation of the <i>Hepsetus</i> from the Congo basin with the revalidation of <i>H. microlepis</i> (Teleostei: Hepsetidae).....	273

Cover photograph

Geophagus diamantinensis (photograph by José L. O. Mattos)
José L. O. Mattos, Wilson J. E. M. Costa and Alexandre C. A. Santos
(this volume pp. 209–220)

Articles appearing in this journal are indexed in:

AQUATIC SCIENCES and FISHERIES ABSTRACTS
BIOLIS - BIOLOGISCHE LITERATUR INFORMATION SENCKENBERG
CAMBRIDGE SCIENTIFIC ABSTRACTS
CURRENT CONTENTS/ AGRICULTURE, BIOLOGY & ENVIRONMENTAL SCIENCES and SCIE
FISHLIT
ZOOLOGICAL RECORD